

## SURVEILLANCE REPORT

# Weekly influenza surveillance overview

29 November 2013

## Main surveillance developments in week 47/2013 (18–24 November 2013)

*This first page contains the main developments for this week and can be printed separately or together with the more detailed information that follows.*

For week 47/2013:

- All 28 reporting countries recorded low-intensity influenza activity.
- Of 357 sentinel specimens tested across 23 countries, three were positive for influenza A virus.
- Three hospitalised laboratory-confirmed influenza cases were reported by the UK.

Since the start of the 2013–2014 influenza surveillance period in week 40/2013, there has been no evidence of sustained influenza activity in Europe.

**Sentinel surveillance of influenza-like illness (ILI)/ acute respiratory infection (ARI):** Low intensity was recorded by all 28 reporting countries. For more information, [click here](#).

**Virological surveillance:** Twenty-three countries collected and tested 357 sentinel specimens, of which three (0.8%) were positive for influenza A virus. For more information, [click here](#).

**Hospital surveillance of laboratory-confirmed influenza cases.** The UK reported three hospitalised patients with laboratory-confirmed influenza: two with influenza type A, one with type B infection. For more information, [click here](#).

# Sentinel surveillance (ILI/ARI)

## Weekly analysis – epidemiology


For week 47/2013, clinical data were reported by 28 countries, all of which recorded low-intensity influenza activity, the lowest category of reporting (Table 1, Map1).

Geographic patterns of influenza activity were reported as sporadic by Denmark, France, Ireland, Norway, Spain, Sweden and the UK (Scotland). All other countries reported no activity (Table 1, Map 2).

Increasing trends were reported by Bulgaria, Estonia, Poland and the UK (Northern Ireland), while all other countries reported stable trends (Table 1, Map 2). The incidence of ILI/ARI was below epidemic thresholds in all countries.

**Map 1. Intensity for week 47/2013****Intensity**

- [Grey square] No report
- [Yellow square] Low
- [Orange square] Medium
- [Red square] High
- [Dark red square] Very High



Liechtenstein

Luxembourg

Malta

(C) ECDC/Dundas/TESSy


\* A type/subtype is reported as dominant when at least ten samples have been detected as influenza positive in the country and of those > 40 % are positive for the type/subtype.

Legend:

<b>No report</b>	Intensity level was not reported	+	Increasing clinical activity
<b>Low</b>	No influenza activity or influenza at baseline levels	-	Decreasing clinical activity
<b>Medium</b>	Usual levels of influenza activity	=	Stable clinical activity
<b>High</b>	Higher than usual levels of influenza activity		
<b>Very high</b>	Particularly severe levels of influenza activity		

**Map 2. Geographic spread for week 47/2013****Geographic spread**

- [Grey square] No Report
- [Light yellow square] No Activity
- [Yellow square] Sporadic
- [Orange square] Local
- [Red square] Regional
- [Dark red square] Widespread



Liechtenstein

Luxembourg

Malta

(C) ECDC/Dundas/TESSy

\* A type/subtype is reported as dominant when at least ten samples have been detected as influenza positive in the country and of those > 40 % are positive for the type/subtype.

**Legend:**

<b>No report</b>	Activity level was not reported	+	Increasing clinical activity
<b>No activity</b>	No evidence of influenza virus activity (clinical activity remains at baseline levels)	-	Decreasing clinical activity
<b>Sporadic</b>	Isolated cases of laboratory confirmed influenza infection	=	Stable clinical activity
<b>Local outbreak</b>	Increased influenza activity in local areas (e.g. a city within a region, or outbreaks in two or more institutions (e.g. schools) within a region (laboratory confirmed)		
<b>Regional activity</b>	Influenza activity above baseline levels in one or more regions with a population comprising less than 50% of the country's total population (laboratory confirmed)		
<b>Widespread</b>	Influenza activity above baseline levels in one or more regions with a population comprising 50% or more of the country's population (laboratory confirmed)		

**Table 1.** Epidemiological and virological overview by country, week 47/2013

Country	Intensity	Geographic spread	Trend	No. of sentinel specimens	Dominant type	Percentage positive	ILI per 100 000	ARI per 100 000	Epidemiological overview	Virological overview
Austria	Low	No activity	Stable	2	None	0.0	833.2	-	Graphs	Graphs
Belgium	Low	No activity	Stable	16	None	0.0	40.5	1831.9	Graphs	Graphs
Bulgaria	Low	No activity	Increasing	3	None	0.0	-	1035.4	Graphs	Graphs
Croatia				0	None	0.0	-	-	Graphs	Graphs
Cyprus	Low	No activity	Stable	-	-	0.0	-*	-*	Graphs	Graphs
Czech Republic	Low	No activity	Stable	-	-	0.0	24.8	854.8	Graphs	Graphs
Denmark	Low	Sporadic	Stable	4	None	0.0	27.4	-	Graphs	Graphs
Estonia	Low	No activity	Increasing	3	None	0.0	7.1	237.4	Graphs	Graphs
Finland	Low	No activity	Stable	8	None	0.0	-	-	Graphs	Graphs
France	Low	Sporadic	Stable	52	None	0.0	-	1531.8	Graphs	Graphs
Germany	Low	No activity	Stable	56	None	0.0	-	1050.2	Graphs	Graphs
Greece	Low	No activity	Stable	5	None	0.0	48.7	-	Graphs	Graphs
Hungary	Low	No activity	Stable	15	None	0.0	64.2	-	Graphs	Graphs
Iceland				0	-	0.0	-	-	Graphs	Graphs
Ireland	Low	Sporadic	Stable	8	None	12.5	4.2	-	Graphs	Graphs
Italy	Low	No activity	Stable	10	None	0.0	94.1	-	Graphs	Graphs
Latvia	Low	No activity	Stable	0	None	0.0	0.0	813.7	Graphs	Graphs
Lithuania	Low	No activity	Stable	9	None	0.0	0.9	504.1	Graphs	Graphs
Luxembourg	Low	No activity	Stable	6	-	0.0	-*	-*	Graphs	Graphs
Malta	Low	No activity	Stable	0	None	0.0	-*	-*	Graphs	Graphs
Netherlands	Low	No activity	Stable	8	None	0.0	33.9	-	Graphs	Graphs
Norway	Low	Sporadic	Stable	6	None	0.0	22.4	-	Graphs	Graphs
Poland	Low	No activity	Increasing	4	None	0.0	192.9	-	Graphs	Graphs
Portugal	Low	No activity	Stable	0	None	0.0	6.6	-	Graphs	Graphs
Romania	Low	No activity	Stable	2	-	0.0	1.0	636.9	Graphs	Graphs
Slovakia	Low	No activity	Stable	3	None	0.0	129.7	1468.6	Graphs	Graphs
Slovenia	Low	No activity	Stable	5	None	0.0	0.0	830.3	Graphs	Graphs
Spain	Low	Sporadic	Stable	69	None	2.9	15.4	-	Graphs	Graphs
Sweden	Low	Sporadic	Stable	26	None	0.0	6.7	-	Graphs	Graphs
UK - England	Low	No activity	Stable	-	None	0.0	1.6	231.7	Graphs	Graphs
UK - Northern Ireland	Low	No activity	Increasing	4	None	0.0	18.8	385.1	Graphs	Graphs
UK - Scotland	Low	Sporadic	Stable	32	None	0.0	7.8	397.3	Graphs	Graphs
UK - Wales	Low	No activity	Stable	1	None	0.0	6.6	-	Graphs	Graphs
<b>Europe</b>				<b>357</b>		<b>0.8</b>				<b>Graphs</b>

\*Incidence per 100 000 is not calculated for these countries as no population denominator is provided.  
Liechtenstein does not report to the European Influenza Surveillance Network.

## Description of the system

Surveillance is based on nationally organised sentinel networks of physicians, mostly general practitioners (GPs), covering at least 1 to 5% of the population in their countries. All EU/EEA Member States (except Liechtenstein) participate. Depending on their country's choice, each sentinel physician reports the weekly number of patients seen with ILI, ARI, or both to a national focal point. From the national level, both numerator and denominator data are then reported to the European Surveillance System (TESSy) database. Additional semi-quantitative indicators of intensity, geographic spread, and trend of influenza activity at the national level are also reported.

# Virological surveillance

## Weekly analysis – virology

For week 47/2013, 23 countries tested a total of 357 sentinel specimens, of which three (0.8%) were positive for influenza A virus (Tables 1–2, Figures 1–2).

In addition, 23 non-sentinel source specimens (e.g. specimens collected for diagnostic purposes in hospitals) were found to be positive for influenza virus: 16 were type A and seven were type B viruses. The two subtyped influenza A viruses were of A(H1)pdm09 subtype (Table 2).

Of the three antigenic characterisations of influenza viruses reported since week 40/2013, two have been characterised as A(H1)pdm09 A/California/7/2009 (H1N1)-like, and one B(Yamagata) lineage virus could not be attributed to any of the reportable categories (Table 3).

Since week 40/2013, 27 genetic characterisations of influenza viruses have been reported for sentinel and non-sentinel specimens. Of these, 13 clustered with A(H1)pdm09 genetic group 6 represented by A/St Petersburg/27/2011, 12 with A(H3) the A/Perth/16/2009 clade, falling within genetic group 3C, represented by A/Texas/50/2012, and two with B(Yamagata)-lineage clade represented by B(Yamagata)-lineage clade 2 represented by B/Massachusetts/02/2012 (Table 4). For details of the current virus strains recommended by WHO for vaccine preparation [click here](#).

More details on viruses that circulated between 1 January and 31 May 2013 can be found in the [September report](#) prepared by the European Reference Laboratory Network for Human Influenza (ERLI-Net) coordination team.



Since week 40/2013, 17 A(H1)pdm09, eight A(H3) viruses and one B virus have been tested for susceptibility to the neuraminidase inhibitors oseltamivir and zanamivir, but none showed genetic or phenotypic ( $IC_{50}$ ) evidence for reduced inhibition.

In week 47/2013, 12 countries reported 543 respiratory syncytial virus (RSV) detections. RSV detections have increased for the third consecutive week but are still at low levels compared to the same time last season (Figure 3).

**Table 2. Weekly and cumulative influenza virus detections by type, subtype and surveillance system, weeks 40–47/2013**

Virus type/subtype	Current period Sentinel	Current period Non-sentinel	Season Sentinel	Season Non-sentinel
Influenza A	3	16	27	188
A(H1)pdm09	1	2	14	51
A(H3)	1	0	10	32
A(sub-type unknown)	1	14	3	105
Influenza B	0	7	4	47
B(Vic) lineage	0	0	0	0
B(Yam) lineage	0	0	0	7
Unknown lineage	0	7	4	40
<b>Total influenza</b>	<b>3</b>	<b>23</b>	<b>31</b>	<b>235</b>

Note: A(H1)pdm09 and A(H3) include both N-subtyped and non-N-subtyped viruses


**Figure 1. Proportion of sentinel specimens positive for influenza virus, weeks 40–47/2013****Figure 2. Number of sentinel specimens positive for influenza virus, by type, subtype and week of report, weeks 40–47/2013**

**Table 3.** Results of antigenic characterisations of sentinel and non-sentinel influenza virus isolates, weeks 40–47/2013

Antigenetic group	Number of viruses
A(H1)pdm09 A/California/7/2009 (H1N1)-like	2
B(Yam) lineage not attributed to category	1

**Table 4.** Results of genetic characterisations of sentinel and non-sentinel influenza virus isolates, weeks 40–47/2013

Phylogenetic group	Number of viruses
A(H1)pdm09 group 6 representative A/St Petersburg/27/2011	13
A(H3) clade repr. A/Perth/16/2009 – A/Texas/50/2012 subgroup (3C)	12
B(Yam)-lineage clade 2 representative B/Massachusetts/02/2012	2

**Figure 3.** Respiratory syncytial virus (RSV) detections, sentinel and non-sentinel, weeks 40–47/2013 and weeks 40/2012-20/2013.

## Description of the system

According to the nationally defined sampling strategy, sentinel physicians take nasal or pharyngeal swabs from patients with ILI, ARI or both and send the specimens to influenza-specific reference laboratories for virus detection, (sub-)typing, antigenic or genetic characterisation and antiviral susceptibility testing.

For details of the current virus strains recommended by WHO for vaccine preparation [click here](#).

# Hospital surveillance – severe influenza disease

## Weekly analysis of hospitalised laboratory-confirmed influenza cases

Since week 40/2013, three countries have reported 18 hospitalised laboratory-confirmed influenza cases (Table 5).

For week 47/2013, two hospitalised influenza A cases and one influenza B case were reported by the UK (Table 6).

Of the 18 hospitalised laboratory-confirmed influenza cases reported since week 40/2013, ten cases were related to infection with influenza virus type A and eight to infection with influenza virus type B (Table 6).

**Table 5.** Cumulative number of hospitalised laboratory-confirmed influenza cases, weeks 40–47/2013

Country	Number of cases	Incidence of cases per 100 000	Estimated population covered
Ireland	1		
Sweden	1		
United Kingdom	16	0.03	63 705 030
<b>Total</b>	<b>18</b>		

**Table 6.** Number of hospitalised laboratory-confirmed influenza cases by influenza type and subtype, week 47/2013 and cumulative for the season

Pathogen	Number of cases during current week	Cumulative number of cases since the start of the season
Influenza A	2	10
A(H1)pdm09	1	4
A(H3)		
A(sub-typing not performed)	1	6
Influenza B	1	8
<b>Total</b>	<b>3</b>	<b>18</b>

## The EuroMOMO mortality monitoring system

Week 47: All-cause mortality has been within the normal range for all reporting countries. Further details are available on <http://www.euromomo.eu>

*This report was written by an editorial team at the European Centre for Disease Prevention and Control (ECDC): Cornelis Adlrich, Eeva Broberg, Julien Beauté and René Snacken. The bulletin text was reviewed by European Reference Laboratory Network for Human Influenza (ERLI-Net) coordination team: Adam Meijer, Rod Daniels, John McCauley and Maria Zambon. On behalf of the EISN members, the bulletin text was reviewed by Maja Sočan (Inštitut za varovanje zdravja), Allison Waters (University College Dublin) and Tyra Grove Krause (Statens Serum Institut, Copenhagen). In addition, the report is reviewed by experts of WHO Regional Office for Europe.*

*Maps and commentary published in this Weekly Influenza Surveillance Overview (WISO) do not represent a statement on the part of ECDC or its partners on the legal or border status of the countries and territories shown.*

*All data published in the WISO are up-to-date on the day of publication. Past this date, however, published data should not be used for longitudinal comparisons as countries tend to retrospectively update their database.*

© European Centre for Disease Prevention and Control, Stockholm, 2013